

SUPPORT VARIETIES FOR MODULES OVER STACKED MONOMIAL ALGEBRAS

TAKAHIKO FURUYA AND NICOLE SNASHALL

ABSTRACT. Let Λ be a finite-dimensional (D, A) -stacked monomial algebra. In this paper, we give necessary and sufficient conditions for the variety of a simple Λ -module to be nontrivial. This is then used to give structural information on the algebra Λ , as it is shown that if the variety of every simple module is nontrivial, then Λ is a D -Koszul monomial algebra. We also provide examples of (D, A) -stacked monomial algebras which are not self-injective but nevertheless satisfy the finite generation conditions **(Fg1)** and **(Fg2)** of [2], from which we can characterize all modules with trivial variety.

INTRODUCTION

Let K be an algebraically closed field, and let $\Lambda = KQ/I$ be an indecomposable finite-dimensional algebra, where Q is a finite quiver and I is an admissible ideal. Let Λ^e be the enveloping algebra $\Lambda \otimes_K \Lambda^{\text{op}}$. The Hochschild cohomology ring of Λ is defined to be $\text{HH}^*(\Lambda) = \text{Ext}_{\Lambda^e}^*(\Lambda, \Lambda) = \bigoplus_{n \geq 0} \text{Ext}_{\Lambda^e}^n(\Lambda, \Lambda)$ with the Yoneda product. We let $\mathcal{N}$ denote the ideal in $\text{HH}^*(\Lambda)$ which is generated by the homogeneous nilpotent elements.

In this paper we consider support varieties for modules over (D, A) -stacked monomial algebras. This class was introduced by Green and Snashall in [4], and the (D, A) -stacked monomial algebras of infinite global dimension were characterized in [3] as precisely the monomial algebras for which every projective module in a minimal projective resolution of $\Lambda/\mathfrak{r}$ over Λ is generated in a single degree and for which the Ext algebra is finitely generated as a K -algebra (but not finite-dimensional). Note that we write $\mathfrak{r}$ for the Jacobson radical of Λ . The class of (D, A) -stacked monomial algebras includes the Koszul monomial algebras (equivalently, the quadratic monomial algebras) and the D -Koszul algebras of Berger [1].

Support varieties for modules over any finite-dimensional algebra Λ were introduced by Snashall and Solberg in [7], where the support variety $V(M)$ of a finitely generated Λ -module M was defined by

$$V(M) = \{\mathfrak{m} \in \text{MaxSpec HH}^*(\Lambda)/\mathcal{N} \mid \text{Ann}_{\text{HH}^*(\Lambda)} \text{Ext}_{\Lambda}^*(M, M) \subseteq \mathfrak{m}'\}$$

2000 *Mathematics Subject Classification.* Primary: 16E40, 16S37.

Key words and phrases. monomial algebra, D -Koszul, Hochschild cohomology, support variety.

The work in this paper was done while the first author was visiting the University of Leicester in 2008/2009. The first author would like to express his gratitude for the hospitality and support from the University of Leicester. The second author wishes to thank the University of Leicester for granting study leave.

where $\mathbf{m}'$ denotes the inverse image of $\mathbf{m}$ in $\mathrm{HH}^*(\Lambda)$. There is a unique maximal graded ideal $\mathbf{m}_{\mathrm{gr}}$ in $\mathrm{HH}^*(\Lambda)/\mathcal{N}$, and $\{\mathbf{m}_{\mathrm{gr}}\} \subseteq V(M)$ for all non-zero finitely generated Λ -modules M ([7, Proposition 3.4]). The variety of M is then said to be trivial if $V(M) = \{\mathbf{m}_{\mathrm{gr}}\}$.

It is clear that if Λ is of finite global dimension then every homogeneous element of $\mathrm{HH}^*(\Lambda)$ of strictly positive degree is nilpotent, and so $\mathrm{HH}^*(\Lambda)/\mathcal{N} \cong K$. It then follows that every module has trivial variety. Since we are interested in the geometric and algebraic connections between varieties and homological properties of modules, we restrict ourselves to algebras of infinite global dimension.

In this paper we study support varieties for modules over a (D, A) -stacked monomial algebra Λ . In Theorem 2.6, we give necessary and sufficient conditions for a simple Λ -module to have trivial variety. This leads to Theorem 2.10, where we provide new structural information on the algebra Λ , namely, we show that if every simple Λ -module has nontrivial variety then $A = 1$ and so Λ is a D -Koszul algebra. It was shown in [2], that many of the properties of support varieties for group algebras have analogues in the more general case under the assumption that certain finiteness conditions **(Fg1)** and **(Fg2)** hold. In particular, [2, Theorem 2.5] shows that under these conditions, the algebra is necessarily Gorenstein (that is, it has finite injective dimension as both a left and a right module), and moreover, the variety of a module is trivial if and only if the module has finite projective dimension. In Section 3, we give examples of (D, A) -stacked monomial algebras which are not self-injective algebras but nevertheless where conditions **(Fg1)** and **(Fg2)** hold. This gives some new and interesting examples where the properties of support varieties for group algebras have analogues for algebras which are not self-injective.

1. BACKGROUND

We fix the notation of this paper. A finite-dimensional algebra $\Lambda = K\mathcal{Q}/I$ is a monomial algebra if I is generated by a set of paths in $K\mathcal{Q}$ each of length at least 2. We fix a minimal generating set ρ for the ideal I , and refer to an element of ρ as a *relation*. We assume further that K is an algebraically closed field with $\mathrm{char} K \neq 2$, and that Λ is indecomposable of infinite global dimension.

An arrow α starts at the vertex $\mathbf{o}(\alpha)$ and ends at the vertex $\mathbf{t}(\alpha)$; arrows in a path are read from left to right. If $p = \alpha_1\alpha_2 \cdots \alpha_n$ is a path in $K\mathcal{Q}$ where $\alpha_1, \dots, \alpha_n$ are arrows, then we set $\mathbf{o}(p) = \mathbf{o}(\alpha_1)$ and $\mathbf{t}(p) = \mathbf{t}(\alpha_n)$. We denote the *length* of a path p by $\ell(p)$. An arrow α is the *first arrow* (respectively *last arrow*) of a relation r in ρ if $r = \alpha p$ (respectively $r = p\alpha$) for some path p .

The definition of a (D, A) -stacked monomial algebra was given in [4] in terms of sets $\mathcal{R}^n$ which are constructed from overlaps of paths in $\mathcal{Q}$. These sets $\mathcal{R}^n$ are also used to define a minimal projective bimodule resolution of Λ . For more details on overlaps see [4].

Definition 1.1. [4, Definition 3.1] Let $\Lambda = K\mathcal{Q}/I$ be a finite-dimensional monomial algebra, where I has a minimal set of generators ρ . Then Λ is said to be a (D, A) -stacked monomial algebra if there is some $D \geq 2$ and $A \geq 1$ such that, for all $n \geq 2$ and $R^n \in \mathcal{R}^n$,

$$\ell(R^n) = \begin{cases} \frac{n}{2}D & n \text{ even} \\ \frac{(n-1)}{2}D + A & n \text{ odd.} \end{cases}$$

In particular all relations in ρ are of length D .

The structure of the Hochschild cohomology ring modulo nilpotence of a (D, A) -stacked monomial algebra was determined in [4, Theorem 3.4]. We use this result heavily in this paper, and therefore we now recall the notation and the result.

A *closed path* C in $\mathcal{Q}$ at the vertex v is a non-trivial path C in $K\mathcal{Q}$ with $C = vCv$ for some vertex v . If C is a closed path at the vertex v then we say that v is *not internal* to C if $C = v\sigma_1v\sigma_2v$ for paths σ_1, σ_2 implies that $\sigma_1 = v$ or $\sigma_2 = v$.

For $A \geq 1$, a *closed A-trail* T in $\mathcal{Q}$ is a non-trivial closed path $T = \alpha_0\alpha_1 \cdots \alpha_{m-1}$ in $K\mathcal{Q}$ such that $\alpha_0, \dots, \alpha_{m-1}$ are all distinct paths of length A . By setting $T_0 = T$ and

$$\begin{aligned} T_1 &= \alpha_1 \cdots \alpha_{m-1} \alpha_0 \\ T_2 &= \alpha_2 \cdots \alpha_0 \alpha_1 \\ &\vdots \\ T_{m-1} &= \alpha_{m-1} \alpha_0 \cdots \alpha_{m-2}, \end{aligned}$$

then we say that $\{T_0, T_1, \dots, T_{m-1}\}$ is a *complete set of closed A-trails* on the A-trail $T = \alpha_0\alpha_1 \cdots \alpha_{m-1}$.

Now, let $d \geq 2$ and write $d = Nm + l$ where $0 \leq l \leq m - 1$ and $N \geq 0$. For $t \in \mathbb{N}$, let $[t] \in \{0, 1, \dots, m - 1\}$ denote the residue of t modulo m . Let $W = T_0^N \alpha_0 \alpha_1 \cdots \alpha_{l-1}$ with the conventions that if $N = 0$ then $T_0^N = \mathfrak{o}(\alpha_0)$ and if $l = 0$ then $W = T_0^N$. More generally, for $k = 0, 1, \dots, m - 1$, define $\sigma^k(W) = T_k^N \alpha_k \alpha_{k+1} \cdots \alpha_{k+l-1}$ with the conventions that

- (i) if $t \geq m$ then $\alpha_t = \alpha_{[t]}$,
- (ii) if $N = 0$ then $T_k^N = e_k$, and
- (iii) if $l = 0$ then $\sigma^k(W) = T_k^N$.

Define ρ_T to be the set

$$\rho_T = \{W, \sigma(W), \dots, \sigma^{m-1}(W)\}.$$

We say that ρ_T is the *set of paths of length dA that are associated to the A-trail T* . Note that $\{W, \sigma(W), \dots, \sigma^{m-1}(W)\}$ is also the set of paths of length dA that is associated to each A-trail T_k for $k = 0, \dots, m - 1$.

Let Λ be a (D, A) -stacked monomial algebra of infinite global dimension. Then [4, Proposition 3.3] tells us that $D = dA$ for some $d \geq 2$.

Let $C_1, \dots, C_u$ be all the closed paths in the quiver $\mathcal{Q}$ at the vertices $v_1, \dots, v_u$ respectively, such that for each C_i with $1 \leq i \leq u$, we have $C_i \neq p_i^{r_i}$ for any path p_i with $r_i \geq 2$, $C_i^d \in \rho$, and there are no overlaps of C_i^d with any relation in $\rho \setminus \{C_i^d\}$. (Note that it follows that $\ell(C_i) = A$.)

Let $T_{u+1}, \dots, T_r$ be all the distinct closed A-trails in the quiver $\mathcal{Q}$ such that for each T_i with $u + 1 \leq i \leq r$, the set ρ_{T_i} of paths of length $D = dA$ which are associated to the trail T_i is contained in ρ but, if $T_i = \alpha_{i0}\alpha_{i1} \cdots \alpha_{im_i-1}$, then each path α_{ij} of length A has no overlaps with any relation in $\rho \setminus \rho_{T_i}$. (We assume that there is no repetition amongst these closed paths and closed A-trails, that is, $\{C_1, \dots, C_u\} \cap \{T_{u+1}, \dots, T_r\} = \emptyset$.)

Then, [4, Theorem 3.4] proves that

$$\mathrm{HH}^*(\Lambda)/\mathcal{N} \cong K[x_1, \dots, x_r]/\langle x_a x_b \mid a \neq b \rangle$$

where

- (a) for $j = 1, \dots, u$, the vertices $v_1, \dots, v_u$ are distinct, and the element x_j corresponding to the closed path C_j is in degree 2 and is represented by the map $P^2 \rightarrow \Lambda$ where, for $R^2 \in \mathcal{R}^2$,

$$\mathfrak{o}(R^2) \otimes \mathfrak{t}(R^2) \mapsto \begin{cases} v_j & \text{if } R^2 = C_j^d \\ 0 & \text{otherwise,} \end{cases}$$

- (b) and for $j = u + 1, \dots, r$, let $T_{j,0}, \dots, T_{j,m_j-1}$ denote the complete set of A -trails on the closed path T_j . Then the element x_j corresponding to the closed A -trail T_j is, in the above notation, in degree $2\mu_j$ where $\mu_j = m_j / \gcd(d, m_j)$ and is represented by the map $P^{2\mu_j} \rightarrow \Lambda$ where, for $R^{2\mu_j} \in \mathcal{R}^{2\mu_j}$,

$$\mathfrak{o}(R^{2\mu_j}) \otimes \mathfrak{t}(R^{2\mu_j}) \mapsto \begin{cases} \mathfrak{o}(T_{j,k}) & \text{if } R^{2\mu_j} = T_{j,k}^{d/\gcd(d, m_j)} \\ & \text{for } k = 0, \dots, m_j - 1 \\ 0 & \text{otherwise.} \end{cases}$$

Throughout this paper, whenever we refer to a closed path C_i in the quiver $\mathcal{Q}$, for some $1 \leq i \leq u$, then we are assuming without further comment that C_i is a closed path at the vertex v_i , such that $C_i \neq p_i^{r_i}$ for some path p_i with $r_i \geq 2$, $C_i^d \in \rho$ where $d = D/A$, and there are no overlaps of C_i^d with any relation in $\rho \setminus \{C_i^d\}$. Similarly, when we refer to a closed A -trail T_i in the quiver $\mathcal{Q}$, for some $u + 1 \leq i \leq r$, then we are assuming without further comment that the set ρ_{T_i} of paths of length D which are associated to the trail T_i is contained in ρ but, if $T_i = \alpha_{i0}\alpha_{i1} \cdots \alpha_{im_i-1}$, then each path α_{ij} of length A has no overlaps with any relation in $\rho \setminus \rho_{T_i}$.

2. SUPPORT VARIETIES FOR SIMPLE MODULES

In this section we give necessary and sufficient conditions for the variety of a simple module to be nontrivial. We first recall the general fact that if (P, δ) is a minimal projective Λ^e -resolution of Λ , then, for any simple Λ -module S , we have that $(P \otimes_\Lambda S, \delta \otimes_\Lambda S)$ is a minimal projective Λ -resolution of S , and so the complex $\text{Hom}_\Lambda(P \otimes_\Lambda S, S)$ has zero differential.

For each vertex v_j of the quiver $\mathcal{Q}$, for $1 \leq j \leq u$, let S_j be the simple module corresponding to v_j . Let $A_j = \text{Ann}_{\text{HH}^*(\Lambda)} \text{Ext}_\Lambda^*(S_j, S_j)$, so A_j is a graded ideal of $\text{HH}^*(\Lambda)$ and is properly contained in $\text{HH}^*(\Lambda)$. Let C_j be a closed path in $\mathcal{Q}$ at the vertex v_j , for some $1 \leq j \leq u$. Then we say that S_j is *associated to the closed path* C_j . We start by considering simple modules associated to one of the closed paths $C_1, \dots, C_u$.

Lemma 2.1. *With the notation already introduced, fix an integer j with $1 \leq j \leq u$, so that C_j is a closed path in the quiver $\mathcal{Q}$ at the vertex v_j . Let z be a non-zero homogeneous element of $\text{HH}^*(\Lambda)$ of the form $z = \sum_{i=1}^r c_i x_i^{l_i}$ with $l_i \geq 0$, $c_i \in K$ and $c_j \neq 0$. Then z is not contained in A_j .*

Proof. If $\deg z = 0$, then $z = c \cdot 1_{\text{HH}^*(\Lambda)} \in \text{HH}^0(\Lambda) \setminus \text{rad HH}^0(\Lambda)$ for some non-zero element $c \in K$, so $z \notin A_j$.

Now suppose $\deg z > 0$, and assume for contradiction that $z \in A_j$. Then $l_j > 0$. Since C_j is a closed path at the vertex v_j , we know that x_j is in degree 2. Hence $\deg z = 2l_j$. Thus we have $\deg z = 2l_j = 2l_t = 2l_w \mu_w$ for all $1 \leq t \leq u$ and

$u + 1 \leq w \leq r$. We write $|z|$ for $\deg z$. From the proof of [4, Proposition 2.5], for $1 \leq t \leq u$, the element $x_t^{l_t} \in \mathrm{HH}^{|z|}(\Lambda)$ is represented by the map $f_t : P^{|z|} \rightarrow \Lambda$ where, for $R \in \mathcal{R}^{|z|}$,

$$(1) \quad f_t(\mathfrak{o}(R) \otimes \mathfrak{t}(R)) = \begin{cases} v_t & \text{if } R = C_t^{dl_t}, \\ 0 & \text{otherwise,} \end{cases}$$

and from the proof of [4, Proposition 2.9], for $u + 1 \leq w \leq r$, the element $x_w^{l_w} \in \mathrm{HH}^{|z|}(\Lambda)$ is represented by the map $f_w : P^{|z|} \rightarrow \Lambda$ where, for $R \in \mathcal{R}^{|z|}$,

$$(2) \quad f_w(\mathfrak{o}(R) \otimes \mathfrak{t}(R)) = \begin{cases} \mathfrak{o}(T_{w,k}) & \text{if } R = T_{w,k}^{dl_w / \gcd(d, m_w)}, \\ & \text{for } k = 0, \dots, m_w - 1 \\ 0 & \text{otherwise.} \end{cases}$$

Then we have

$$f_i(\mathfrak{o}(C_j^{dl_j}) \otimes \mathfrak{t}(C_j^{dl_j})) = \begin{cases} v_j & \text{if } i = j \\ 0 & \text{otherwise.} \end{cases}$$

Since $z \in A_j$, the element $z \otimes_\Lambda S_j \in \mathrm{Ext}_\Lambda^{|z|}(S_j, S_j)$ is zero. Moreover since $z \otimes_\Lambda S_j$ is represented by $\sum_{i=1}^r (c_i f_i \otimes_\Lambda S_j) : P^{|z|} \otimes_\Lambda S_j \rightarrow \Lambda \otimes_\Lambda S_j$, there is a left Λ -homomorphism $\Phi : P^{|z|-1} \otimes_\Lambda S_j \rightarrow \Lambda \otimes_\Lambda S_j$ such that $\sum_{i=1}^r (c_i f_i \otimes_\Lambda S_j) = \Phi \circ (\delta^{|z|} \otimes_\Lambda S_j)$. So we have the following situation

$$\begin{array}{ccc} P^{|z|} \otimes_\Lambda S_j & \xrightarrow{\delta^{|z|} \otimes_\Lambda S_j} & P^{|z|-1} \otimes_\Lambda S_j \\ \downarrow z \otimes_\Lambda S_j & \nwarrow \Phi & \\ \Lambda \otimes_\Lambda S_j & & \end{array}$$

Then, by the remark at the start of this section, we have $\Phi \circ (\delta^{|z|} \otimes_\Lambda S_j) = 0$. But

$$\begin{aligned} & \left(\sum_{i=1}^r (c_i f_i \otimes_\Lambda S_j) \right) (\mathfrak{o}(C_j^{dl_j}) \otimes \mathfrak{t}(C_j^{dl_j}) \otimes_\Lambda v_j) \\ &= (c_j f_j \otimes_\Lambda S_j) (\mathfrak{o}(C_j^{dl_j}) \otimes \mathfrak{t}(C_j^{dl_j}) \otimes_\Lambda v_j) \\ &= c_j (v_j \otimes_\Lambda v_j) \end{aligned}$$

which is not zero in $\Lambda \otimes_\Lambda S_j$, since $c_j \neq 0$. So $\sum_{i=1}^r (c_i f_i \otimes_\Lambda S_j)$ is not zero. This is a contradiction. Hence it follows that $z \notin A_j$. $\square$

Proposition 2.2. *With the notation already introduced, fix an integer j with $1 \leq j \leq u$, so that C_j is a closed path in the quiver $\mathcal{Q}$ at the vertex v_j . Then the variety of the simple module S_j associated to C_j is nontrivial.*

Proof. We show first that $(A_j + \mathcal{N})/\mathcal{N}$ is an ideal of the subalgebra

$$R_j = K[x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_r] / \langle x_a x_b \mid a \neq b \rangle$$

of $K[x_1, \dots, x_r] / \langle x_a x_b \mid a \neq b \rangle \cong \mathrm{HH}^*(\Lambda)/\mathcal{N}$.

Let $z + \mathcal{N} \in (A_j + \mathcal{N})/\mathcal{N}$, so that $z = y + \chi$ for some $y \in A_j, \chi \in \mathcal{N}$. Suppose first that z is a homogeneous element of $\mathrm{HH}^*(\Lambda)$. Since $\mathrm{HH}^*(\Lambda)/\mathcal{N} \cong K[x_1, \dots, x_r] / \langle x_a x_b \mid a \neq b \rangle$, we can also write $z = \sum_{i=1}^r c_i x_i^{l_i} + \chi_1$ with $l_i \geq 0$, $c_i \in K$ and $\chi_1 \in \mathcal{N}$. Thus $y = \sum_{i=1}^r c_i x_i^{l_i} + \chi_2$ where $\chi_2 = \chi_1 - \chi \in \mathcal{N}$. Since χ_2 is nilpotent, there is some $n \geq 0$ with $\chi_2^n = 0$, and so $(y - \sum_{i=1}^r c_i x_i^{l_i})^n = 0$. Using the graded commutativity of $\mathrm{HH}^*(\Lambda)$, we then have that $(\sum_{i=1}^r c_i x_i^{l_i})^n \in \mathrm{HH}^*(\Lambda)y$.

Since y is in the ideal A_j and $x_a x_b = 0$ for $a \neq b$, this gives $\sum_{i=1}^r c_i^n x_i^{l_i n} = (\sum_{i=1}^r c_i x_i^{l_i})^n \in A_j$.

Now, we may use Lemma 2.1, to show that either $\sum_{i=1}^r c_i^n x_i^{l_i n} = 0$ or $c_j^n = 0$. If $\sum_{i=1}^r c_i^n x_i^{l_i n} = 0$ then $(\sum_{i=1}^r c_i x_i^{l_i})^n = 0$ so $\sum_{i=1}^r c_i x_i^{l_i} \in \mathcal{N}$. Thus $z \in \mathcal{N}$ and so $z + \mathcal{N} = 0 + \mathcal{N} \in R_j$. On the other hand, if $c_j^n = 0$ then $c_j = 0$ and so $z + \mathcal{N} = \sum_{i=1, i \neq j}^r c_i x_i^{l_i} + \mathcal{N} \in R_j$. In both cases we have that $z + \mathcal{N} \in R_j$.

Now suppose that z is not a homogeneous element of $\mathrm{HH}^*(\Lambda)$. Since A_j and $\mathcal{N}$ are both graded ideals, we have $z = \sum_{i=1}^N z_i$ for some homogeneous elements $z_i = y_i + \chi_i$ with $y_i \in A_j, \chi_i \in \mathcal{N}$ and some $N \geq 0$. From above, we know that $z_i + \mathcal{N} \in R_j$ for $1 \leq i \leq N$, and so $z + \mathcal{N} = \sum_{i=1}^N (z_i + \mathcal{N}) \in R_j$.

Thus $(A_j + \mathcal{N})/\mathcal{N}$ is contained in R_j . Therefore $(A_j + \mathcal{N})/\mathcal{N}$ is contained in the maximal ideal $(x_1, \dots, x_{j-1}, x_j - \alpha, x_{j+1}, \dots, x_r)/\langle x_a x_b \mid a \neq b \rangle$ of $\mathrm{HH}^*(\Lambda)/\mathcal{N}$ for all $\alpha \in K$.

Now,

$$\begin{aligned} V(S_j) &= \{\mathfrak{m} \in \mathrm{MaxSpec} \mathrm{HH}^*(\Lambda)/\mathcal{N} \mid A_j \subseteq \mathfrak{m}'\} \\ &= \{\mathfrak{m} \in \mathrm{MaxSpec} \mathrm{HH}^*(\Lambda)/\mathcal{N} \mid (A_j + \mathcal{N})/\mathcal{N} \subseteq \mathfrak{m}\}. \end{aligned}$$

Thus $(x_1, \dots, x_{j-1}, x_j - \alpha, x_{j+1}, \dots, x_r)/\langle x_a x_b \mid a \neq b \rangle \in V(S_j)$ for all $\alpha \in K$. Hence the variety of S_j is nontrivial. $\square$

Let T_j be a closed A -trail in the quiver $\mathcal{Q}$ for some $u+1 \leq j \leq r$. For k with $0 \leq k \leq m_{j-1}$, we denote by S_{jk} the simple module corresponding to the vertex $\mathfrak{o}(T_{j,k})$. We say that the simple modules S_{jk} , for $0 \leq k \leq m_{j-1}$, are *associated to the closed A -trail T_j* . Let $A_{jk} = \mathrm{Ann}_{\mathrm{HH}^*(\Lambda)} \mathrm{Ext}_{\Lambda}^*(S_{jk}, S_{jk})$, so A_{jk} is a graded ideal of $\mathrm{HH}^*(\Lambda)$ and is properly contained in $\mathrm{HH}^*(\Lambda)$. We now show the analogue of Lemma 2.1 for closed A -trails.

Lemma 2.3. *Keeping our notation, fix integers j and k with $u+1 \leq j \leq r$ and $0 \leq k \leq m_j - 1$, so that T_j is a closed A -trail in the quiver $\mathcal{Q}$, and $T_{j,0}, \dots, T_{j,m_j-1}$ is the complete set of A -trails on T_j . Let z be a non-zero homogeneous element of $\mathrm{HH}^*(\Lambda)$ of the form $z = \sum_{i=1}^r c_i x_i^{l_i}$ with $l_i \geq 0$, $c_i \in K$ and $c_j \neq 0$. Then z is not contained in A_{jk} .*

Proof. If $\deg z = 0$, then $z = c \cdot 1_{\mathrm{HH}^*(\Lambda)} \in \mathrm{HH}^0(\Lambda) \setminus \mathrm{rad} \mathrm{HH}^0(\Lambda)$ for some non-zero element $c \in K$, so $z \notin A_{jk}$.

Now suppose that $\deg z > 0$, and assume for contradiction that $z \in A_{jk}$. Then $l_j > 0$. Since T_j is a closed A -trail, we know that x_j is in degree $2\mu_j$. Hence $\deg z = 2l_j \mu_j$. Thus we have $\deg z = 2l_j \mu_j = 2l_t = 2l_w \mu_w$ for all $1 \leq t \leq u$ and $u+1 \leq w \leq r$. Again we denote $\deg z$ by $|z|$. Recall from the proof of Lemma 2.1 that, for $1 \leq t \leq u$, the element $x_t^{l_t} \in \mathrm{HH}^{|z|}(\Lambda)$ is represented by the map $f_t : P^{|z|} \rightarrow \Lambda$ given by (1), and, for $u+1 \leq w \leq r$, the element $x_w^{l_w} \in \mathrm{HH}^{|z|}(\Lambda)$ is represented by the map $f_w : P^{|z|} \rightarrow \Lambda$ given by (2). Then we have

$$f_i(\mathfrak{o}(T_{j,k}^{dl_j/\gcd(d,m_j)}) \otimes \mathfrak{o}(T_{j,k}^{dl_j/\gcd(d,m_j)})) = \begin{cases} \mathfrak{o}(T_{j,k}) & \text{if } i = j \\ 0 & \text{otherwise.} \end{cases}$$

Since $z \in A_{jk}$, the element $z \otimes_{\Lambda} S_{jk} \in \mathrm{Ext}_{\Lambda}^{|z|}(S_{jk}, S_{jk})$ is zero. Moreover since $z \otimes_{\Lambda} S_{jk}$ is represented by $\sum_{i=1}^r (c_i f_i \otimes_{\Lambda} S_{jk}) : P^{|z|} \otimes_{\Lambda} S_{jk} \rightarrow \Lambda \otimes_{\Lambda} S_{jk}$, there is a left Λ -homomorphism $\Phi : P^{|z|-1} \otimes_{\Lambda} S_{jk} \rightarrow \Lambda \otimes_{\Lambda} S_{jk}$ such that $\sum_{i=1}^r (c_i f_i \otimes_{\Lambda}$

$S_{jk}) = \Phi \circ (\delta^{|z|} \otimes_{\Lambda} S_{jk})$. Then, by the remark at the start of this section, we have $\Phi \circ (\delta^{|z|} \otimes_{\Lambda} S_{jk}) = 0$. But

$$\begin{aligned} & \left(\sum_{i=1}^r (c_i f_i \otimes_{\Lambda} S_{jk}) \right) (\mathfrak{o}(T_{j,k}^{dl_j / \gcd(d, m_j)}) \otimes \mathfrak{o}(T_{j,k}^{dl_j / \gcd(d, m_j)}) \otimes_{\Lambda} \mathfrak{o}(T_{j,k})) \\ &= (c_j f_j \otimes_{\Lambda} S_{jk}) (\mathfrak{o}(T_{j,k}^{dl_j / \gcd(d, m_j)}) \otimes \mathfrak{o}(T_{j,k}^{dl_j / \gcd(d, m_j)}) \otimes_{\Lambda} \mathfrak{o}(T_{j,k})) \\ &= c_j (\mathfrak{o}(T_{j,k}) \otimes_{\Lambda} \mathfrak{o}(T_{j,k})), \end{aligned}$$

which is not zero in $\Lambda \otimes_{\Lambda} S_{jk}$, since $c_j \neq 0$. So $\sum_{i=1}^r (c_i f_i \otimes_{\Lambda} S_{jk})$ is not zero. This is a contradiction. Hence it follows that $z \notin A_{jk}$. $\square$

Now, in a similar way to Proposition 2.2, we have the following result concerning the varieties of simple modules associated to closed A -trails.

Proposition 2.4. *Keeping our notation, fix integers j and k with $u+1 \leq j \leq r$ and $0 \leq k \leq m_j - 1$, so that T_j is a closed A -trail in the quiver $\mathcal{Q}$, and $T_{j,0}, \dots, T_{j,m_j-1}$ is the complete set of A -trails on T_j . Then the variety of the simple module S_{jk} is nontrivial.*

Proof. First we show that $(A_{jk} + \mathcal{N})/\mathcal{N}$ is an ideal of the subalgebra

$$R_j = K[x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_r] / \langle x_a x_b \mid a \neq b \rangle$$

of $K[x_1, \dots, x_r] / \langle x_a x_b \mid a \neq b \rangle \cong \text{HH}^*(\Lambda)/\mathcal{N}$.

Let $z + \mathcal{N} \in (A_{jk} + \mathcal{N})/\mathcal{N}$. A similar argument to that used in the proof of Proposition 2.2 together with Lemma 2.3 shows that $z + \mathcal{N} \in R_j$. Thus $(A_{jk} + \mathcal{N})/\mathcal{N}$ is contained in R_j . Hence $(A_{jk} + \mathcal{N})/\mathcal{N}$ is contained in the maximal ideal $(x_1, \dots, x_{j-1}, x_j - \alpha, x_{j+1}, \dots, x_r) / \langle x_a x_b \mid a \neq b \rangle$ of $\text{HH}^*(\Lambda)/\mathcal{N}$ for all $\alpha \in K$.

Again following Proposition 2.2, we have that the variety of S_{jk} is nontrivial. $\square$

Now we consider the varieties of modules which are not associated to one of the closed paths $C_1, \dots, C_u$ or the closed A -trails $T_{u+1}, \dots, T_r$ in the quiver $\mathcal{Q}$.

Proposition 2.5. *Let S be a simple module which is not associated to one of the closed paths $C_1, \dots, C_u$ or the closed A -trails $T_{u+1}, \dots, T_r$ in the quiver $\mathcal{Q}$. Then the variety of S is trivial.*

Proof. If $\text{HH}^*(\Lambda)/\mathcal{N} \cong K$, then the variety of every finitely generated module is trivial. In particular, the variety of each simple module is trivial.

Assume now that $\text{HH}^*(\Lambda)/\mathcal{N} \not\cong K$, so we have some $r \geq 1$ with

$$\text{HH}^*(\Lambda)/\mathcal{N} \cong K[x_1, \dots, x_r] / \langle x_a x_b \mid a \neq b \rangle.$$

Recall that, for $1 \leq i \leq u$, the element x_i corresponding to the closed path C_i at the vertex v_i is represented by the map $P^2 \rightarrow \Lambda$ where, for $R^2 \in \mathcal{R}^2$,

$$\mathfrak{o}(R^2) \otimes \mathfrak{t}(R^2) \mapsto \begin{cases} v_i & \text{if } R^2 = C_i^d \\ 0 & \text{otherwise,} \end{cases}$$

and, for $u+1 \leq j \leq r$, the element x_j corresponding to the closed A -trail T_j is represented by the map $P^{2\mu_j} \rightarrow \Lambda$ where, for $R^{2\mu_j} \in \mathcal{R}^{2\mu_j}$,

$$\mathfrak{o}(R^{2\mu_j}) \otimes \mathfrak{t}(R^{2\mu_j}) \mapsto \begin{cases} \mathfrak{o}(T_{j,k}) & \text{if } R^{2\mu_j} = T_{j,k}^{d / \gcd(d, m_j)} \\ & \text{for } k = 0, \dots, m_j - 1 \\ 0 & \text{otherwise.} \end{cases}$$


Since S is not associated to one of the closed paths C_i at the vertex v_i for $1 \leq i \leq u$ or to one of the closed A -trails T_j for $u+1 \leq j \leq r$, it follows that $v_i \otimes_\Lambda S = 0$ for $1 \leq i \leq u$, and $\mathfrak{o}(T_{j,k}) \otimes_\Lambda S = 0$ for $u+1 \leq j \leq r$ and $0 \leq k \leq m_j - 1$. Thus the map $x_j \otimes_\Lambda S$ is zero for all $1 \leq j \leq r$. So $(x_1, \dots, x_r) \subseteq \text{Ann}_{\text{HH}^*(\Lambda)} \text{Ext}_\Lambda^*(S, S)$. Hence if $\mathfrak{m} \in \text{MaxSpec HH}^*(\Lambda)$ with $\text{Ann}_{\text{HH}^*(\Lambda)} \text{Ext}_\Lambda^*(S, S) \subseteq \mathfrak{m}$ then $\mathfrak{m} = (x_1, \dots, x_r)$. Therefore $V(S) = \{(x_1, \dots, x_r)\}$, so that the variety of S is trivial. $\square$

From Propositions 2.2, 2.4, and 2.5, we have the following theorem.

Theorem 2.6. *Let S be a simple module. Then the variety of S is trivial if and only if S is not associated to one of the closed paths $C_1, \dots, C_u$ or to one of the closed A -trails $T_{u+1}, \dots, T_r$ in the quiver $\mathcal{Q}$.*


We now give some examples to illustrate this theorem.

Example 2.7. Let $\Lambda = K\mathcal{Q}/I$ where $\mathcal{Q}$ is the quiver




and I is the ideal $\langle \alpha\beta, \beta\gamma, \gamma\delta, \delta\alpha \rangle$. Then the simple modules S_1, S_2, S_3, S_4 of Λ corresponding to the vertices 1, 2, 3, 4 respectively are associated to the closed 1-trail $\alpha\beta\gamma\delta$. So, by Theorem 2.6, the varieties of S_1, S_2, S_3, S_4 are nontrivial, whereas the varieties of the simple modules corresponding to the vertices 5, 6 are trivial.

Example 2.8. Let $\Lambda = K\mathcal{Q}/I$ where $\mathcal{Q}$ is the quiver



and $I = \langle \alpha\beta, \beta\gamma, \gamma\alpha, \zeta\eta, \eta\theta, \theta\zeta \rangle$. Then all simple modules of Λ are associated to closed A -trails in $\mathcal{Q}$. In fact, the simple module corresponding to the vertex 1 is associated to both the closed 1-trails $\alpha\beta\gamma$ and $\zeta\eta\theta$. Thus, by Theorem 2.6, the varieties of all simple modules of Λ are nontrivial.

Example 2.9. Let $\Lambda = K\mathcal{Q}/I$ where $\mathcal{Q}$ is the quiver



and $I = \langle \alpha\beta\gamma\delta\alpha\beta, \gamma\delta\alpha\beta\gamma\delta \rangle$. Then the simple modules S_1, S_3 corresponding to the vertices 1, 3 are associated to the closed 2-trail $\alpha\beta\gamma\delta$. Hence, by Theorem 2.6, the varieties of S_1, S_3 are nontrivial, whereas the varieties of the simple modules corresponding to the vertices 2, 4 are trivial.

We complete this section with the following theorem which uses information on the varieties of the simple modules to give structural information on Λ .

Theorem 2.10. *Let $\Lambda = K\mathcal{Q}/I$ be a finite-dimensional (D, A) -stacked monomial algebra. Suppose that each simple module has nontrivial variety. Then $A = 1$. Thus Λ is a D -Koszul monomial algebra.*

Proof. There is a simple Λ -module with nontrivial variety, so $\mathrm{HH}^*(\Lambda)/\mathcal{N} \not\cong K$. Thus, from [4, Theorem 3.4], there exists an integer $r \geq 1$ with

$$\mathrm{HH}^*(\Lambda)/\mathcal{N} \cong K[x_1, \dots, x_r]/\langle x_a x_b \mid a \neq b \rangle,$$

where $x_1, \dots, x_u$ correspond to the closed trails $C_1, \dots, C_u$ and $x_{u+1}, \dots, x_r$ correspond to the distinct closed A -trails $T_{u+1}, \dots, T_r$. From Theorem 2.6, every simple right Λ -module is associated to a closed path $C_1, \dots, C_u$ or a closed A -trail $T_{u+1}, \dots, T_r$. Thus, for each vertex v , we may fix a relation R_v^2 in $\{C_1^d, \dots, C_u^d\} \cup (\cup_{i=u+1}^r \rho_{T_i})$ so that $R_v^2 = vR_v^2$.

Choose a vertex $v_1 \in \mathcal{Q}$. Then $v_1 = \mathfrak{o}(R_{v_1}^2)$ for some $R_{v_1}^2$ as above. Denote the first arrow in $R_{v_1}^2$ by a_1 , and let $v_2 = \mathfrak{t}(a_1)$. Then $v_2 = \mathfrak{o}(R_{v_2}^2)$. Again, let a_2 denote the first arrow in $R_{v_2}^2$ and let $v_3 = \mathfrak{t}(a_2)$. Continuing in this way gives a path $a_1 a_2 a_3 \dots$ in $K\mathcal{Q}$. Since $\mathcal{Q}$ has a finite number of vertices, eventually some vertex v_i must repeat, so that there is a closed path $a_1 a_2 a_3 \dots a_p$ in $K\mathcal{Q}$ with $\mathfrak{o}(a_1) = \mathfrak{t}(a_p)$ and $p \geq 1$.

The algebra Λ is finite-dimensional, so there is a subpath of $(a_1 a_2 \dots a_p)^N$ for $N \gg 0$ which lies in ρ , and so is necessarily of length D . Denote this element of ρ by R_*^2 and, by relabelling if necessary, suppose that the last arrow of R_*^2 is a_1 , so that $R_*^2 = a_j \dots a_{j+D-2} a_1$ for some j . Thus we have the situation:

$$\begin{array}{c} \overbrace{a_j \dots a_{j+D-2} \overbrace{a_1}^{R_{v_1}^2}}^{R_*^2} \end{array}$$

Now $R_{v_1}^2$ overlaps R_*^2 . But $R_{v_1}^2 \in \{C_1^d, \dots, C_u^d\} \cup (\cup_{i=u+1}^r \rho_{T_i})$.

If $R_{v_1}^2 = C_i^d$ for some $1 \leq i \leq u$, then $R_*^2 = C_i^d$, since we know that there are no overlaps of C_i^d with any other relation in ρ . But then, the first arrow of R_*^2 equals the first arrow of $R_{v_1}^2$, so $a_j = a_1$ and $R_*^2 = a_1 a_{j+1} \dots a_{j+D-2} a_1 = C_i^d$. But [4, Lemma 2.2] gives us that $\mathfrak{o}(R_*^2) = \mathfrak{o}(a_1)$ is not internal to C_i , so we must have $R_*^2 = a_1^D$ and $C_i = a_1$. Thus $A = \ell(C_i) = \ell(a_1) = 1$.

Otherwise, we have $R_{v_1}^2 \in \rho_{T_i}$ for some $u+1 \leq i \leq r$. Since there are no overlaps of relations in ρ_{T_i} with relations in $\rho \setminus \rho_{T_i}$, we know $R_*^2 \in \rho_{T_i}$ as well. By the construction of ρ_{T_i} , we know that there is a relation $\tilde{R}_*^2$ in ρ_{T_i} which ends at $\mathfrak{o}(R_{v_1}^2) = v_1$. Then R_*^2 and $\tilde{R}_*^2$ lie in ρ_{T_i} , and R_*^2 overlaps $\tilde{R}_*^2$ with overlap of length $D+1$. This overlap is necessarily maximal and may be illustrated as follows.

$$\begin{array}{c} \overbrace{\overbrace{a_1}^{R_{v_1}^2}}^{\tilde{R}_*^2} \overbrace{\quad}^{R_*^2} \end{array}$$

Hence there is an element of $\mathcal{R}^3$ of length $D+1$. Thus $D+1 = D+A$, so $A = 1$. $\square$

3. CONDITIONS (Fg1) AND (Fg2)

In this section we give some examples of (D, A) -stacked monomial algebras which are not self-injective but nevertheless satisfy the following two finiteness conditions found in [2]:

(Fg1) There is a graded subalgebra H of $\mathrm{HH}^*(\Lambda)$ such that

- (i) H is a commutative Noetherian ring.
- (ii) $H^0 = \mathrm{HH}^0(\Lambda) = Z(\Lambda)$.

(Fg2) The Ext algebra of Λ


$$E(\Lambda) = \mathrm{Ext}_{\Lambda}^*(\Lambda/\mathfrak{r}, \Lambda/\mathfrak{r}) = \bigoplus_{i \geq 0} \mathrm{Ext}_{\Lambda}^i(\Lambda/\mathfrak{r}, \Lambda/\mathfrak{r})$$

is a finitely generated H -module.

In the case where these conditions hold, we may apply the results of [2] to give further significant information on support varieties for Λ -modules. In particular, we can apply [2, Theorem 2.5] to show that, for a finite-dimensional K -algebra Λ satisfying **(Fg1)** and **(Fg2)**, the algebra Λ is necessarily Gorenstein, and, for a finitely generated Λ -module M , the variety of M is trivial if and only if the projective dimension of M is finite.

We now present some examples of (D, A) -stacked monomial algebras of infinite global dimension which are not self-injective but which illustrate the possible behaviours with respect to the conditions **(Fg1)** and **(Fg2)**. In the first two examples, conditions **(Fg1)** and **(Fg2)** hold.

Example 3.1. As in Example 2.8, let $\Lambda = K\mathcal{Q}/I$ where $\mathcal{Q}$ is the quiver



and $I = \langle \alpha\beta, \beta\gamma, \gamma\alpha, \zeta\eta, \eta\theta, \theta\zeta \rangle$. Then, Λ is a Koszul monomial algebra (so is a $(2, 1)$ -stacked monomial algebra), and the variety of every simple module is non-trivial.

Now, we claim that Λ satisfies **(Fg1)** and **(Fg2)**. By [4, Theorem 3.4], $\mathrm{HH}^*(\Lambda)/\mathcal{N}$ is isomorphic to the subalgebra $K[x, y]/(xy)$ of $\mathrm{HH}^*(\Lambda)$, where x corresponds to the 1-trail $\alpha\beta\gamma$ and is represented by the map $P^6 \rightarrow \Lambda$ where, for $R^6 \in \mathcal{R}^6 = \{(\alpha\beta\gamma)^2, (\beta\gamma\alpha)^2, (\gamma\alpha\beta)^2, (\zeta\eta\theta)^2, (\eta\theta\zeta)^2, (\theta\zeta\eta)^2\}$,

$$\mathfrak{o}(R^6) \otimes \mathfrak{t}(R^6) \mapsto \begin{cases} e_1 & \text{if } R^6 = (\alpha\beta\gamma)^2, \\ e_2 & \text{if } R^6 = (\beta\gamma\alpha)^2, \\ e_3 & \text{if } R^6 = (\gamma\alpha\beta)^2, \\ 0 & \text{otherwise,} \end{cases}$$

and y corresponds to the 1-trail $\zeta\eta\theta$ and is represented by the map $P^6 \rightarrow \Lambda$ where, for $R^6 \in \mathcal{R}^6$,

$$\mathfrak{o}(R^6) \otimes \mathfrak{t}(R^6) \mapsto \begin{cases} e_1 & \text{if } R^6 = (\zeta\eta\theta)^2, \\ e_4 & \text{if } R^6 = (\eta\theta\zeta)^2, \\ e_5 & \text{if } R^6 = (\theta\zeta\eta)^2, \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to verify that $\mathrm{HH}^0(\Lambda) = Z(\Lambda) = K$. Let H be the graded subalgebra of $\mathrm{HH}^*(\Lambda)$ generated by 1, x and y . Then it follows that $H \cong K[x, y]/(xy)$ and $H^0 = K$. Furthermore, H is a commutative Noetherian ring. Thus Λ satisfies **(Fg1)**.

Now, we have $\mathcal{R}^0 = \{e_1, \dots, e_5\}$ and $\mathcal{R}^1 = \{\alpha, \beta, \gamma, \zeta, \eta, \theta\}$. Since Λ is a Koszul monomial algebra, we know that $E(\Lambda)$ is generated by $\mathcal{R}^0$ and $\mathcal{R}^1$ as an algebra.

Let n be a positive integer, and write $n = 6q + r$ with $0 \leq r \leq 5$. For $i = 1, 2, 3$, let $P_{i,r}$ be the path of length r which lies on the closed 1-trail $\alpha\beta\gamma$ such that $\mathfrak{o}(P_i) = i$. Similarly, for $i = 1, 4, 5$, let $Q_{i,r}$ be the path of length r which lies on the closed 1-trail $\zeta\eta\theta$ such that $\mathfrak{o}(Q_i) = i$. Thus both $P_{i,r}$ and $Q_{j,r}$ represent elements of degree r in $E(\Lambda)$ for $i = 1, 2, 3$, and $j = 1, 4, 5$. Moreover, since $(\alpha\beta\gamma)^2$, $(\beta\gamma\alpha)^2$ and $(\gamma\alpha\beta)^2$ represent elements of degree 6 in $E(\Lambda)$, it follows that $(\alpha\beta\gamma)^{2q}$, $(\beta\gamma\alpha)^{2q}$ and $(\gamma\alpha\beta)^{2q}$ represent elements of degree $6q$ in $E(\Lambda)$ for $n = 6q + r$ with $0 \leq r \leq 5$, where $(\alpha\beta\gamma)^0 = e_1$, $(\beta\gamma\alpha)^0 = e_2$ and $(\gamma\alpha\beta)^0 = e_3$. By definition of $\mathcal{R}^n$, we get

$$\mathcal{R}^n = \{(\alpha\beta\gamma)^{2q}P_{1,r}, (\beta\gamma\alpha)^{2q}P_{2,r}, (\gamma\alpha\beta)^{2q}P_{3,r}, (\zeta\eta\theta)^{2q}Q_{1,r}, (\eta\theta\zeta)^{2q}Q_{4,r}, (\theta\zeta\eta)^{2q}Q_{5,r}\}$$

for $n = 6q + r$ with $0 \leq r \leq 5$. Also, we know from Section 1 of [5] that the set $\cup_{i \geq 0} \mathcal{R}^i$ is a multiplicative basis of $E(\Lambda)$. Now the action of H on $E(\Lambda)$ is induced by the ring homomorphism $H \rightarrow E(\Lambda)$ determined by

$$\begin{cases} 1 & \mapsto 1, \\ x & \mapsto (\alpha\beta\gamma)^2 + (\beta\gamma\alpha)^2 + (\gamma\alpha\beta)^2, \\ y & \mapsto (\zeta\eta\theta)^2 + (\eta\theta\zeta)^2 + (\theta\zeta\eta)^2, \end{cases}$$

so we have $\mathcal{R}^n = \{x^q P_{1,r}, x^q P_{2,r}, x^q P_{3,r}, y^q Q_{1,r}, y^q Q_{4,r}, y^q Q_{5,r}\}$ for $n = 6q + r$ with $0 \leq r \leq 5$. This shows that $E(\Lambda)$ is generated by $\cup_{0 \leq i \leq 5} \mathcal{R}^i$ as a left H -module, and hence is a finitely generated left H -module. Hence Λ satisfies **(Fg2)**.

Moreover $e_1\Lambda$ is not injective, so Λ is not self-injective. We know from [2, Theorem 2.5] that Λ is necessarily Gorenstein, and it is easy to verify that the injective dimension of Λ is 1. Thus we have given an example of a nonself-injective Koszul algebra where every simple module has nontrivial variety and **(Fg1)** and **(Fg2)** hold.

Example 3.2. As in Example 2.9, let $\Lambda = K\mathcal{Q}/I$ where $\mathcal{Q}$ is the quiver

$$\begin{array}{ccc} 1 & \xrightarrow{\alpha} & 2 \\ \delta \uparrow & & \downarrow \beta \\ 4 & \xleftarrow{\gamma} & 3 \end{array}$$

and $I = \langle \alpha\beta\gamma\delta\alpha\beta, \gamma\delta\alpha\beta\gamma\delta \rangle$. Then the varieties of the simple modules corresponding to 1, 3 are nontrivial, whereas the varieties of the simple modules corresponding to 2, 4 are trivial. Thus, for a simple module S , we see directly that the variety of S is trivial if and only if the projective dimension of S is finite. Moreover $e_3\Lambda$ is not injective, so that Λ is not self-injective.

We now show that Λ satisfies **(Fg1)** and **(Fg2)**, and hence Λ must be Gorenstein by [2, Theorem 2.5]. By [4, Theorem 3.4], $\mathrm{HH}^*(\Lambda)/\mathcal{N} \cong K[x]$ where x corresponds to the 2-trail $\alpha\beta\gamma\delta$ and is represented by the map $P^4 \rightarrow \Lambda$, where, for $R^4 \in \mathcal{R}^4 = \{(\alpha\beta\gamma\delta)^3, (\gamma\delta\alpha\beta)^3\}$,

$$\mathfrak{o}(R^4) \otimes \mathfrak{t}(R^4) \mapsto \begin{cases} e_1 & \text{if } R^4 = (\alpha\beta\gamma\delta)^3, \\ e_3 & \text{if } R^4 = (\gamma\delta\alpha\beta)^3. \end{cases}$$

Let $z = \alpha\beta\gamma\delta + \beta\gamma\delta\alpha + \gamma\delta\alpha\beta + \delta\alpha\beta\gamma \in \Lambda$. Then it is easy to see that $z \in Z(\Lambda)$ and $\mathrm{HH}^0(\Lambda) = Z(\Lambda) = K[z]/(z^2)$. Moreover, let H be the subalgebra of $\mathrm{HH}^*(\Lambda)$ generated by 1, z , x . Then it follows that $H^0 = K[z]/(z^2)$ and $H \cong K[x, z]/(z^2)$, and hence H is a commutative Noetherian ring. Therefore Λ satisfies **(Fg1)**.

Now, we have $\mathcal{R}^0 = \{e_1, e_2, e_3, e_4\}$ and $\mathcal{R}^1 = \{\alpha, \beta, \gamma, \delta\}$. Let

$$R_1 = \alpha\beta\gamma\delta\alpha\beta, \quad R_2 = \gamma\delta\alpha\beta\gamma\delta, \quad R'_1 = \alpha\beta\gamma\delta\alpha\beta\gamma\delta, \quad R'_2 = \gamma\delta\alpha\beta\gamma\delta\alpha\beta.$$

Then, for $i = 1, 2$, R_i and R'_i represent elements of degrees 2 and 3 of $E(\Lambda)$ respectively, and we have $\mathcal{R}^2 = \{R_1, R_2\}$ and $\mathcal{R}^3 = \{R'_1, R'_2\}$. We know from [3] that $E(\Lambda)$ is generated by $\mathcal{R}^0$, $\mathcal{R}^1$, $\mathcal{R}^2$, and $\mathcal{R}^3$ as an algebra. Let n be an integer with $n \geq 4$, and write $n = 4q + r$ with $0 \leq r \leq 3$ (so $q \geq 1$). Then, we get

$$\mathcal{R}^n = \begin{cases} \{(R_1 R_2)^q, (R_2 R_1)^q\} & \text{if } r = 0, \\ \{(R_1 R_2)^{q-1} R_1 R'_2, (R_2 R_1)^{q-1} R_2 R'_1\} & \text{if } r = 1, \\ \{(R_1 R_2)^q R_1, (R_2 R_1)^q R_2\} & \text{if } r = 2, \\ \{(R_1 R_2)^q R'_1, (R_2 R_1)^q R'_2\} & \text{if } r = 3. \end{cases}$$

Also, we know from Section 1 of [5] that the set $\cup_{i \geq 0} \mathcal{R}^i$ is a multiplicative basis of $E(\Lambda)$. Now the action of H on $E(\Lambda)$ is induced by the ring homomorphism $H \rightarrow E(\Lambda)$ determined by

$$\begin{cases} 1 & \mapsto 1, \\ x & \mapsto R_1 R_2 + R_2 R_1, \\ z & \mapsto 0, \end{cases}$$

so that, for $n \geq 4$, we have

$$\mathcal{R}^n = \begin{cases} \{x^q e_1, x^q e_2\} & \text{if } r = 0, \\ \{x^{q-1} R_1 R'_2, x^{q-1} R_2 R'_1\} & \text{if } r = 1, \\ \{x^q R_1, x^q R_2\} & \text{if } r = 2, \\ \{x^q R'_1, x^q R'_2\} & \text{if } r = 3. \end{cases}$$

This shows that $E(\Lambda)$ is generated by $\cup_{0 \leq i \leq 5} \mathcal{R}^i$ as a left H -module. So Λ satisfies **(Fg2)**.


Moreover it can be seen that the injective dimension of Λ is 2. In fact, $e_2 \Lambda$ and $e_4 \Lambda$ are injective modules, and moreover we have the following injective resolutions of $e_1 \Lambda$ and $e_3 \Lambda$ respectively:

$$\begin{aligned} 0 \rightarrow e_1 \Lambda &\longrightarrow I(2) \longrightarrow I(4) \longrightarrow I(3) \rightarrow 0, \\ 0 \rightarrow e_3 \Lambda &\longrightarrow I(4) \longrightarrow I(2) \longrightarrow I(1) \rightarrow 0, \end{aligned}$$

where $I(j)$ is the indecomposable injective module corresponding to the vertex j for $1 \leq j \leq 4$. Thus the injective dimensions of $e_1 \Lambda$ and $e_3 \Lambda$ are 2, and so the injective dimension of Λ is 2. Thus we have given an example of a nonself-injective monomial algebra where some simple modules have trivial variety, some simple modules have nontrivial variety, and **(Fg1)** and **(Fg2)** hold.

In the next two examples, conditions **(Fg1)** and **(Fg2)** do not hold.

Example 3.3. Let $\mathcal{Q}$ be the quiver



and let $\Lambda = K\mathcal{Q}/I$ where $I = \langle \alpha^2, \beta\gamma, \gamma\beta, ab, ac, ba, ca \rangle$. Then Λ is Koszul (so is a $(2, 1)$ -stacked monomial algebra). Also, the vertex 1 is associated to the closed path α , and the vertices 2 and 3 are associated to the 1-trail $\beta\gamma$. So, by Theorem 2.6, the variety of each simple module is nontrivial.

Now, there is the following minimal projective resolution of the indecomposable injective right Λ -module $I(3)$ corresponding to the vertex 3:


$$\cdots \longrightarrow U_3 \longrightarrow U_2 \longrightarrow U_1 \longrightarrow e_3\Lambda \oplus e_3\Lambda \longrightarrow I(3) \rightarrow 0,$$

where, for $i \geq 1$,

$$U_i = \begin{cases} (e_1\Lambda)^{\oplus 3 \cdot 2^{m-1}} & \text{if } i = 2m - 1, \\ (e_2\Lambda)^{\oplus 3 \cdot 2^{m-1}} & \text{if } i = 2m. \end{cases}$$

Hence, the projective dimension of $I(3)$ is infinite, which implies that the injective dimension of Λe_3 is infinite. So Λ is not Gorenstein. Hence, by [2, Theorem 2.5], there is no H satisfying **(Fg1)** and **(Fg2)**. Therefore we have given an example of a nonself-injective Koszul monomial algebra where the variety of every simple module is nontrivial, and **(Fg1)** and **(Fg2)** do not hold.

Example 3.4. Let $\mathcal{Q}$ be the quiver



and let $\Lambda = K\mathcal{Q}/I$ where $I = \langle \alpha\beta, \alpha\gamma, \gamma\alpha, \beta\alpha \rangle$. Then Λ is a Koszul monomial algebra with radical squared zero. From [4, Theorem 3.4], $\mathrm{HH}^*(\Lambda)/\mathcal{N} \cong K$, and so the variety of every finitely generated module is trivial. Thus all simple modules of Λ have trivial variety. Moreover, it is easy to see that all the simple Λ -modules have infinite projective dimension. It now follows from [2, Theorem 2.5] that there is no H such that **(Fg1)** and **(Fg2)** hold. In particular, Λ is not Gorenstein. Thus we have given an example of a nonself-injective Koszul monomial algebra where the variety of every simple module is trivial, and **(Fg1)** and **(Fg2)** do not hold.

REFERENCES

- [1] R. Berger, *Koszulity for nonquadratic algebras*, J. Algebra **239** (2001), 705–734.
- [2] K. Erdmann, M. Holloway, N. Snashall, Ø. Solberg, R. Taillefer, *Support varieties for selfinjective algebras*, K-Theory **33** (2004), 67–87.
- [3] E. L. Green and N. Snashall, *Finite generation of Ext for a generalization of D-Koszul algebras*, J. Algebra **295** (2006), 458–472.
- [4] E. L. Green and N. Snashall, *The Hochschild cohomology ring modulo nilpotence of a stacked monomial algebra*, Colloq. Math. **105** (2006), 233–258.
- [5] E. L. Green and D. Zacharia, *The cohomology ring of a monomial algebra*, Manuscripta Math. **85** (1994), 11–23.
- [6] D. Happel, *Hochschild cohomology of finite-dimensional algebras*, Séminaire d'Algèbre Paul Dubreil et Marie-Paul Malliavin, (ed. M.-P. Malliavin), Lecture Notes in Mathematics **1404**, Springer, New York, 1989, 108–126.
- [7] N. Snashall and Ø. Solberg, *Support varieties and Hochschild cohomology rings*, Proc. London Math. Soc. (3) **88** (2004), 705–732.

TAKAHIKO FURUYA, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LEICESTER, UNIVERSITY ROAD, LEICESTER, LE1 7RH, ENGLAND
E-mail address: `tf47@mcs.le.ac.uk`

NICOLE SNASHALL, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LEICESTER, UNIVERSITY ROAD, LEICESTER, LE1 7RH, ENGLAND
E-mail address: `N.Snashall@mcs.le.ac.uk`